

MICROBIAL CONTROL OF MOSQUITOES IN WEST GERMANY AND THE HUBEI PROVINCE OF THE PEOPLE'S REPUBLIC OF CHINA

NORBERT BECKER¹ AND XU BOZHAO²

¹*Kommunale Aktionsgemeinschaft zur Bekämpfung der Stechmückenplage e.V.,
6700 Ludwigshafen am Rhein, Europlatz 5, West Germany*

²*Institute of Parasitic Diseases, Hubei Academy of Medical Science,
Wuhan, Hubei, People's Republic of China*

ABSTRACT

In West Germany, along major river systems such as the Upper Rhine, the nuisance caused by flood water mosquitoes very often diminish the quality of life of the local residents. In the Province of Hubei in the People's Republic of China mosquitoes are vectors of malaria, Japanese-B- Encephalitis, and lymphatic filariasis. Due to the incurring resistance, and high development costs, as well as the environmental problems caused by common and newly developed insecticides, the control of these mosquitoes has become increasingly more difficult.

In both parts of the world, microbial agents such as *Bacillus thuringiensis israelensis* and *Bacillus sphaericus* have allowed an effective, cost efficient, and environmentally safe control of these mosquitoes. In West Germany the mosquito population has been reduced about 90% per year by applying approximately 5-10 tons of *B.t.i.*, depending on the fluctuation of the Rhine River water level, in about 500 km² of floodlands. In Hubei, each year about 12 000 hectares of rice fields and sewage ditches are treated with B.t.-187 (local strain) and B.s.C3-41 (local strain) preparations within an integrated control program against *Anopheles sinensis* and *Culex pipiens quinquefasciatus*. The incidence of malaria has thereby been reduced from 5.6 in 1986 to 0.8/10,000 people in 1989.

I INTRODUCTION

Mosquitoes are worldwide the most common vectors of dangerous diseases such as malaria, arbovirus diseases, or filariasis. According to WHO estimates, more than one hundred million cases of malaria occur worldwide each year. More than 2.5 billion people live in areas where malaria is highly endemic.

For example in the Province of Hubei more than 20 million people on both sides of the Yangtsekiang River are threatened by the malaria agent *Plasmodium vivax*. During 1985 more than 50,000 cases of malaria were reported, which corresponds to a yearly incidence of 10/10,000 people. The major vector is the fever mosquito *Anopheles sinensis*, the control of which has recently become more and more difficult with the common insecticides, because of developing resistance as well as ecological and toxicological risks. Other diseases transmitted by mosquitoes are Japanese-B-Encephalitis, a virus disease transmitted by *Culex pipiens quinquefasciatus*, *Cx. tritaeniorhynchus*, *Aedes albopictus*, and *Anopheles sinensis*, and also Brugian and Bancroftian filariasis, transmitted mainly by *An. sinensis* and *Cx. pipiens quinquefasciatus*.

Owing to their massive occurrence along the larger river systems in Europe, mosquitoes, especially floodwater mosquitoes, can be a terrible nuisance. Due to the frequent fluctuations in the water level on the Upper Rhine Valley, mosquito populations there were often extremely dense. The dominant mosquito species are the floodwater mosquitoes *Aedes vexans*, *Ae. sticticus* and

Ae. rossicus, which breed in temporary water bodies resulting from rises in the water level of the Rhine River (Becker, 1989). Because of the nuisance caused by mosquito bites, the quality of life in the affected areas was often extremely reduced. During late afternoons and evenings, or on sultry days, the public was unable to spend any length of time outdoors. It was not uncommon for many hundreds of mosquitoes to attack a pedestrian in just a few minutes.

Culex pipiens molestus is the main pest in houses. Their breeding sites are usually rainwater containers near the houses and other water bodies occurring in summer.

2. CONVENTIONAL MOSQUITO CONTROL

For more than 40 years, chemical insecticides have allowed a relatively simple and economical control of almost all disease vectors. According to WHO calculations, in the early 1980's more than 50.000 t of chemical insecticides were applied yearly in those developing countries especially affected by vector-borne diseases. Even today, DDT is the most widely used insecticide, representing approximately 50% of all insecticides used in the tropics. It was first believed that chemical insecticides alone could eradicate diseases such as malaria or onchocerciasis. However, many vectors soon became increasingly resistant to insecticides. New insecticides had to be developed at high costs, whereby the rapid development of new resistance could not be excluded. Besides the resistance problem, environmental problems such as accumulation in the food web and unselective effects, killing the predators of the mosquitoes as well, accompany the common insecticides. For these reasons the search for environmentally safe, biological methods of insect control was begun. Among all of the microbial control agents for mosquitoes and black flies, the spore-forming bacteria such as *Bacillus thuringiensis israelensis* and *Bacillus sphaericus* are the most promising. This shall be illustrated by two samples.

3. MICROBIAL CONTROL OF MOSQUITOES IN GERMANY

As a reaction to the nuisance caused by mosquitoes, approximately 100 towns and villages on both sides of the Rhine River joined together to form a voluntary mosquito control organization (KABS). The territory of this organization now covers approximately 300 river kilometers (Becker and Ludwig, 1983).

The objective of this program is the reduction of the mosquito population to a tolerable level without damaging the ecologically sensitive riverbank areas and their biocoenosis. This objective has been achieved by the widespread use of microbial agents.

One of the strongest assets of the organization is the well-trained local field staff. There are usually about 300 people active and the yearly budget is about 1 mill. U.S. dollars. Depending on the size of the breeding sites, in every community there are between two and eight people responsible for mosquito control. Besides the community workers, biology students familiar with the local area are employed during flood periods as temporary workers in the control of *Aedes* species, receiving a loan of 6 U.S. dollars.

On the one hand, an integrated control, applying different but compatible methods, reduces the development of resistance. On the other hand, it allows a more pest-specific control. It is important that, from the available methods, always those methods are chosen which are not only effective in the given situation but are also environmentally safe.

The following methods are applied in the mosquito control program in West Germany:

a) Microbiological Methods: For the last eight years, *B.t.i.*-preparations have been mostly used in routine control measures. Today the mosquitoes in more than 90% of the area exclusively are controlled by the KABS with *B.t.i.*. *Bacillus sphaericus* preparations are increasingly being used against *Cx. pipiens molestus*.

b) Water Management and Related Measures: The protection and encouragement of all natural

predators is a very important maxim of our integrated methods. As the predators are not affected by microbial agents, they can continue to feed upon newly hatching mosquito larvae after the breeding sites have been treated. This also has economical consequences, since in most cases no further treatments are then necessary.

c) Surface film method: Only when *B.t.i.* is no longer effective, is a self-spreading biodegradable surface film, which is a mixture of soybean, lecithin, and paraffin applied in concentrations of 0.4–0.8 ml/m² against late fourth instars and pupae.

In 1979 we started testing various available commercial *B.t.i.* products, such as BACTIMOS, TEKNAR, and VECTOBAC (Tables 1 and 2).

As a result of these tests, in routine treatments against first and second instar larvae or in shallow breeding sites, 250 g of *B.t.i.* powder (activity: 6000 AAU/mg) is mixed with 10 liters of screen-filtered pond water per hectare. The mixture is applied by field workers with a hand-held, high-pressure sprayer. On the other hand, in deeper breeding sites or when third or fourth instar larvae are present, 500 g of *B.t.i.* WP is used. This corresponds to about 1.8×10^9 AA International Toxic Units per hectare when about 300 g WP are applied.

In order to achieve a sufficiently high mortality rate under field conditions, one liter of fluid concentrate (Teknar HP-D) or Vectobac 12 AS) is mixed with 9 liters of screen-filtered pond water per hectare.

When high water levels on the Rhine cause widespread inundation or when dense vegetation occurs, *B.t.i.* granules formulated by ourselves are applied. This formulation consists of 50 kg of *B.t.i.* wettable powder (activity: 6000 AAU/mg) in a cement mixer. This is enough to treat about 2 hectares of breeding site either by helicopter, equipped with a SIMPLEX sprayer, or by hand.

Between 1981 and 1989, more than 40,000 hectares of mosquito breeding sites were successfully treated with about 18 tons of *B.t.i.* wettable powder and 17,000 liters of *B.t.i.* flowable concentrate. The *B.t.i.* WP was also used to produce approximately 300 tons of *B.t.i.* sand granules.

The control of *Culex pipiens molestus* near houses is based on providing information to the general public on the biology of the house mosquitoes and strategies for their control. People are asked to destroy all unnecessary water bodies near their homes, to empty the rain water containers at least once a week, or to cover them thoroughly. Fish can also be used as predators or *B.t.i.* briquets can be applied in rainwater containers, the latter allowing efficient control for several weeks. For the last 5 years, we provided about 300,000 *B.t.i.* briquets to the public. Since 1989 we have been testing briquets on the basis of *Bacillus sphaericus* to optimize the *Culex pipiens* control.

As a result of all these measures, the mosquito population in the Upper Rhine Valley was reduced each year by over 90%. This has been highly respected by the public, especially regarding the low costs and the tremendous increase in the quality of life.

4. MICROBIAL CONTROL OF MOSQUITOES IN HUBEI PROVINCE

In a cooperative program between the Province of Hubei and Baden-Württemberg (West Germany), the scientific and organizational prerequisites for the large-scale application of microbial agents are being improved. In detail, the aims of this program are:

- a) Intensive exchange of practical and scientific knowledge as well as technology in the field of integrated mosquito control between West Germany and the Province of Hubei.
- b) Detailed investigations on the biology and ecology of the most important mosquito vectors of human diseases, as a basis for their successful control by microbiological means.
- c) Development of low-cost and environmentally safe control measures, which can be applied at the community level.
- d) Effective improvement of the measures and techniques for routine control operations in both countries in a long-lasting cooperation between the two working groups.

TABLE 1
Field evaluation of various wettable powder formulations of *Bacillus thuringiensis* H-14 against larvae of *Aedes vexans* and *Aedes cantians*

Product	Treatment rate	Habitat surface/depth	Species instars	No. of larvae/10 dips pre- and post-treatment		
				Pretreatment	1	2 (days)
BACTIMOS WP (6000 AAU/mg)	0.1 kg/ha	Swampy woodlands 19.5m ² /0.15m	<i>Ae. cantians</i> (L ₃ -L ₄)	109	22 (79.8)	0 (100)
TEKNAR WP (5800 AAU/mg)	0.1 kg/ha	Swampy woodlands 18.2m ² /0.15m	<i>Ae. cantians</i> (L ₃ -L ₄)	57	43 (24.6)	5 (91.2)
VECTOBAC WP (3600 ITU/mg)	0.1 kg/ha	Swampy woodlands 16m ² /0.1m	<i>Ae. cantians</i> (L ₃ -L ₄)	131	119 (9.2)	67 (48.9)
BACTIMOS WP	0.2 kg/ha	Swampy woodlands 18m ² /0.1m	<i>Ae. cantians</i> (L ₃ -L ₄)	62	3 (95.2)	0 (100)
TEKNAR	0.2 kg/ha	Swampy woodlands 19.5m ² /0.15m	<i>Ae. cantians</i> (L ₃ -L ₄)	81	38 (53.1)	3 (96.3)
VECTOBAC	0.2 kg/ha	Swampy woodlands 24m ² /0.15m	<i>Ae. cantians</i> (L ₃ -L ₄)	106	76 (28.3)	1 (99.1)
TEKNAR WP	0.1 kg/ha	Floodlands 120m ² /0.2m	<i>Ae. vexans</i> (L ₂ -L ₄)	130	95 (26.9)	34 (73.9)
TEKNAR WP	0.2 kg/ha	Floodlands 120m ² /0.2m	<i>Ae. vexans</i> (L ₁ -L ₄)	260	2 (99.2)	0 (100)
BACTIMOS WP	0.2 kg/ha	Floodlands 80/m ² /0.2m	<i>Ae. vexans</i> (L ₁ -L ₄)	759	14 (98.2)	2 (99.7)

() = percent reduction; L₁-L₃ = larval instars

TABLE 2
Field evaluation of various flowable concentrates of *Bacillus thuringiensis H-14* against larvae of *Aedes vexans* and *Aedes cantans*

Product	Treatment rate	Habitat surface/depth	Species instars	Pretreatment	No. of larvae/10 dips pre- and post-treatment			
					1	2	3	4 (days)
TEKNAR 402 SC (1500 AAU/mg)	1 l/ha	Floodlands 225m ² /0.2m	<i>Ae. vexans</i> (L ₁ -L ₄)	386	7 (98.2)	1 (99.7)	0 (100)	0 (100)
TEKNAR 402 SC (1500 AAU/mg)	1 l/ha	Swampy woodlands 6.5m ² /0.2m	<i>Ae. cantans</i> (L ₂ -L ₃)	100	62 (38)	31 (69)	8 (92)	0 (100)
TEKNAR HP-D (3000 AAU/mg)	1 l/ha	Swampy woodlands 6.5m ² /0.2m	<i>Ae. cantans</i> (L ₂ -L ₃)	186	2 (98.9)	0 (100)	0 (100)	0 (100)
TEKNAR HP-D (3000 AAU/mg)	1 l/ha	Swampy woodlands 12m ² /0.3m	<i>Ae. cantans</i> (L ₃)	312	53 (83.1)	16 (94.9)	6 (98.1)	1 (99.7)
VECTOBAC-AS (600 ITU/mg)	1 l/ha	Swampy woodlands 11.2m ² /0.3m	<i>Ae. cantans</i> (L ₃)	328	241 (26.5)	51 (84.5)	44 (86.4)	5 (98.5)

() = percent reduction; L₁-L₄ = larval instars

4.1. Laboratory tests with *Bacillus thuringiensis* and *B.s.* preparations

During the first phase of the program, two *B.t.i.* powder formulations and four *B.s.* preparations were tested in the laboratory of the Institute of Parasitic Diseases in Wuhan (Hubei). The tests were carried out against larvae of *Cx. pipiens quinquefasciatus*, *An. sinensis*, and *Ae. albopictus* at concentrations of 0.001; 0.01; 0.1; 1 and 10 ppm. While the *B.t.i.* preparations were very effective against larvae of *Ae. albopictus* (LC50 = 0.025–0.0568 ppm), they were less active against *Cx. quinquefasciatus* larvae (LC50 = 0.037–0.28 ppm).

On the other hand, a relatively low dosage of *Bacillus sphaericus* was already enough to kill *Culex* larvae (LC50 = 0.001–0.009 ppm), whereas *Aedes* larvae are less sensitive (LC50 = 0.54–0.72 ppm). However, both microbial preparations did not provide satisfactory control of larvae of *An. sinensis* (LC50 = 1.9–10 ppm).

4.2. Field tests with *B.t.i.* and *B.s.*

The most effective laboratory preparations were tested against *Cx. pipiens quinquefasciatus* and *An. sinensis* in eight drainage canals and ponds.

The large scale field tests against *Cx. pipiens quinquefasciatus* larvae with the most effective preparations yielded highly sufficient results at concentrations of 0.5–1 ppm for about one week.

Besides testing for efficiency, we could also show that microbial agents have no negative impact to the ecosystem, especially to the predators of mosquitoes. These field tests, as well as the laboratory experiments, demonstrated that for the control of *Anopheles* larvae, more effective microbial preparations need to be developed in order to reduce the excessive control costs caused by high dosages and repeated treatments. In this context it was of interest to develop a slow releasing floating formulation for the control of *Anopheles* larvae as a "surface feeder."

During the last years, about 10 tons of *B.t.i.*-187 (local strain) and about 14 tons of *B.s.* C3-41 (local strain) have been produced each year by utilization of natural resources in Hubei Province, which was enough to treat about 12,000 hectares of mosquito breeding sites. In 1989 about 200 kg of new floating granules were used against *Anopheles sinensis* larvae treating an area of about 300 hectares of rice fields.

As a result of these measures the incidence of malaria decreased from 5.6 cases/10,000 people in 1986 to 0.8 cases/10,000 people in 1989 (Xu, Becker and Xianqi, 1990).

B.t.i. and *B.s.* preparations offer not only high efficiency against mosquitoes coupled with low treatment costs, but also complete safety for all non-target organisms, which can be promising agents in the fight against dangerous diseases such as malaria and filariasis. It can be assumed that, through applying these microbial agents in an integrated program, at least for an acceptable time-period, the onset of resistance can be prevented.

REFERENCES

- Becker, N. and Ludwig, H.W. 1983. Mosquito control in West Germany. *Bull. Soc. Vector Ecol.* 8(2):85–93.
 Becker, N. 1989. Life strategies of mosquitoes as an adaptation to their habitats. *Bull. Soc. Vector Ecol.* 14(1):6–25.
 Xu, B.Z., Becker, N. and Xianqi, X. 1990. Microbial control of mosquitoes in Hubei Province. (in press)

EVALUATION OF DIVERSE FORMULATIONS OF *BACILLUS THURINGIENSIS*
VAR. *ISRAELENSIS* AGAINST *ANOPHELES ALBIMANUS* IN HONDURAS¹

M.J. PERICH, L.R. BOOBAR, J.C. STIVERS² AND L.A. RIVERA³
U.S. Army Biomedical Research and Development Laboratory,
Fort Detrick, Frederick, MD 21701-5010, U.S.A.

ABSTRACT

Four biorational larvicide formulations, Duplex[®] (methoprene combined with *B.t.i.*), Teknar[®] (*B.t.i.*), Arosurf[®] MSF (Monomolecular Surface Film), and Arosurf MSF combined with Teknar, were evaluated against naturally occurring populations of *Anopheles albimanus* Wiedemann in Honduras. All formulations reduced the mean number of larvae per sample area to 0 within 48 hr posttreatment, and gave significant ($P \leq 0.05$) control through 240 hr posttreatment. The potential for large scale use of these formulations for vector suppression in Honduras is discussed.

INTRODUCTION

Malaria is the primary arthropod-borne communicable disease in Honduras, with over 30,000 cases reported in 1985 (PAHO, 1986). The disease is a major deterrent of socioeconomic development in the tropical Americas. Historically, vector control has been the major component of any malaria control program in endemic areas (Sloff, 1987).

Anopheles albimanus Wiedemann is a major vector of malaria in Central America (Clyde, 1987). This species has become physiologically resistant to many of the conventional insecticides (Brown, 1986). In addition, behavioral modifications (exophagic behavior) in *An. albimanus*'s blood feeding activity, has made standard domiciliary spraying with residual insecticides ineffective (J.C. Stivers, personal communication). These factors, along with the potential insult to the environment by synthetic organic insecticides (Matsumura, 1975), has created the need for biorational alternative control strategies.

Larviciding with biorational formulations has been proposed as one alternative control strategy against malaria vectors as part of an integrated vector control program (Sloff, 1987). Various commercial formulations of *Bacillus thuringiensis* var. *israelensis* (*B.t.i.*) alone, and combined with a monomolecular surface film (Perich et al., 1987) or methoprene (Perich et al., 1988), have been found to be efficacious under laboratory conditions against *An. albimanus*. The purpose of this study was to evaluate these biorational larvicide formulations against *An. albimanus* in various breeding habitats in the Comayagua Valley of Honduras.

¹The opinions or assertions contained herein are the private views of the author(s) and are not to be construed as official or as reflecting the views of the Department of the Army or the Department of Defense.

²Agency for International Development, Tegucigalpa, DC, Honduras, CA.

³Division Control de Vectores, Ministry of Public Health, Tegucigalpa, DC, Honduras, CA.

MATERIALS AND METHODS

The treatment areas were located within a 3 km radius of the city of Comayagua, in the Comayagua Valley, the principal agricultural area of southcentral Honduras. This region, historically an endemic malaria area, contains many drainage/irrigation ditches, small ponds (<0.5 ha), rice fields, and slow moving streams, known sites of *An. albimanus* breeding (Breeland, 1972; Rozeboom, 1941). Treatment applications were randomly assigned to these sites.

The larvicide formulations evaluations were: Teknar[®], a commercial monomolecular organic surface film; Arosurf[®] MSF combined with Teknar (4:1), a noncommercially available formulation; and Duplex[®], a commercially available formulation of Teknar and methoprene combined. All larvicide formulations were applied as technical grade solutions using a 7.6 liter hand-pumped compression sprayer. A separate sprayer was used with each formulation. The application rates in the study were 1.17 liters/ha for the Teknar and Duplex larvicides, and 4.67 liters/ha for the Arosurf MSF larvicide. A homogeneous suspension of the combined formulation of Arosurf MSF and Teknar was ensured by vigorous hand shaking throughout the spray operation.

Sites were selected based on pretreatment sampling of previously known positive *An. albimanus* breeding sites (L.A. Rivera, personal communication). Once a site was verified to contain *An. albimanus* larvae, three, 400 ml dipper samples were taken at three sample points, at a minimum of 5 m apart, prior to treatment. Sample points were marked to ensure that the same points would be sampled posttreatment. Treatments were applied with a fan spray to the sampling point and surrounding water for a 100 m radius in all directions using a sweeping pattern. Posttreatment samples (3 dips/sample point) were taken at 24, 48, 72 and 240 hr.

The mean number of larvae at each posttreatment sampling interval for all formulations was used as the criteria in evaluating their efficacy against *An. albimanus*. A completely randomized experimental design was used; the data were analyzed by an analysis of variance (ANOVA [SAS/STAT, 1985]). The means of the pre- and posttreatment samples were separated by use of Duncan's (1955) multiple range test ($P \leq 0.05$).

RESULTS AND DISCUSSION

All formulations evaluated reduced the number of *An. albimanus* larvae (1st–4th instar) significantly within 24 hr posttreatment (Tables 1 and 2). No statistical difference ($P \leq 0.05$) was found between any two of the formulations. Each formulation provided significant reduction in the number of *An. albimanus* 1st–4th instar larvae through 240 hr posttreatment. Ramoska et al. (1982), van Essen and Hembree (1982), Margalit and Bobroglio (1984), and Ohana et al. (1987) have shown that large amounts of organic matter in the water significantly reduce the field efficacy and persistence of *B.t.i.*. Since organic matter in the treatment area water was relatively low, the high level of initial control and persistence provided in this study was expected.

At 24 hr posttreatment, the Arosurf MSF alone did not provide 100% reduction in the 1st and 2nd larval instar populations as did the other three formulations although 100% mortality was obtained at 48 hr (Table 1). This lower sensitivity of an early *An. albimanus* instars to Arosurf MSF corroborates previous findings described by Perich et al. (1987). Although no significant pupal data were collected in this study (only one pupa for entire test period), the pupicidal activity of Arosurf MSF can be a significant component in the suppression of *An. albimanus* (Perich et al., 1987).

The combined formulation of Arosurf MSF + Teknar has been shown to produce a joint larvicidal/pupicidal action against *An. albimanus* (Perich et al., 1987). This additive effect along with the induced spreading action afforded by Arosurf MSF (Levy et al., 1984), makes this formulation highly efficacious against *An. albimanus*, as shown in Tables 1 and 2. A major operational advantage of using combined formulations (Arosurf MSF + Teknar, Duplex) rather than single component larvicides is that they can be used against a broader range of immature mosquitoes (1st

TABLE 1
Mean number of *Anopheles albimanus* 1st and 2nd larval instar after application
of larvicide formulations in Comayagua, Honduras^{a,b}

Larvicide Formulation	0 hr Pre-trt	24 hrs Post-tr	48 hrs Post-tr	72 hrs Post-tr	240 hrs Post-tr
Teknar	23.7 a	0 a	0 a	0 a	1.5 a
Arosurf MSF	11.7 a	1.0 a	0 a	0 a	0 a
Arosurf MSF + Teknar	11.0 a	0 a	0 a	0 a	0 a
Duplex	17.3 a	0 a	0 a	0 a	1.0 a
Control	18.0 a	16.0 b	12.0 b	17.7 b	11.7 b

^aMean number based on three, 400 ml dipper samples per three sample points per sample area.

^bMeans within a column followed by the same letter are not significantly different ($P \leq 0.05$; Duncan's [1955] multiple range test).

larval instar-pupal stage). A broad, efficacious application range against *An. albimanus* immatures is necessary because of their rapid stadia and continuous breeding in the tropics (Breeland, 1974; Del Carmen et al., 1984).

Certain limiting factors are cosmopolitan for any large scale use of *B.t.i.* and/or monomolecular organic surface film formulations for mosquito control. These limiting factors are: short field persistence by either formulation in areas treated; *B.t.i.* formulations (excluding combined formulation with Arosurf MSF) not remaining suspended in treated water; little canopy penetration when aerially applied; and high cost of treatment. These same factors are then potential influences on the efficacy of all formulations evaluated in this study, when used in large area treatments against *An. albimanus* throughout its geographical distribution.

Short persistence in mosquito breeding habitats is a major disadvantage for both *B.t.i.* (Margalit and Dean, 1985) and monomolecular organic surface films (Levy et al., 1981). In a subsequent control operation in the Comayagua Valley, all formulations were found to provide significant control for only 12 days posttreatment. This limited persistence by *B.t.i.* can, in part, be attributed to its ready adsorption to organic particles (Ohana et al., 1987) and gravity which prevent it from remaining suspended in the water and therefore available for larval ingestion.

TABLE 2
Mean number of *Anopheles albimanus* 3rd and 4th larval instars after application
of larvicide formulations in Comayagua, Honduras^{a,b}

Larvicide Formulation	0 hr Pre-trt	24 hrs Post-tr	48 hrs Post-tr	72 hrs Post-tr	240 hrs Post-tr
Teknar	1.8 a	0 a	0 a	0 a	0 a
Arosurf MSF	1.3 a	0 a	0 a	0 a	0 a
Arosurf MSF + Teknar	1.3 a	0 a	0 a	0 a	0.3 a
Duplex	1.3 a	0 a	0 a	0 a	0.3 a
Control	5.3 b	5.0 b	3.3 b	2.3 b	2.0 b

^aMean number based on three, 400 ml dipper samples per three sample points per sample area.

^bMeans within a column followed by the same letter are not significantly different ($P \leq 0.05$; Duncan's [1955] multiple range test).

Canopy penetration was not a factor in this study because applications were made at ground level ≤ 30 cm from the water surface. Canopy penetration becomes a limiting factor for these formulations when aerial applications are used in treating large areas. Thus, it becomes necessary to add suitable canopy penetrating carriers to the active ingredients (*B.t.i.*, Arosurf MSF, and methoprene) in order to get through the dense vegetation associated with *An. albimanus* larval habitat (Elliott, 1969).

The cost of using any of the formulations is high (Table 3) when considering the potential need for retreatment at 10–12 day intervals. Such high application costs make large area treatment with these formulations prohibitive for most areas in Honduras at the present time.

TABLE 3
Approximate costs for the amount of four larvicide formulations necessary to treat one hectare

Larvicide formulation	Cost per ha (US\$)
Teknar	6.18
Arosurf MSF	22.66
Arosurf MSF + Teknar	28.84
Duplex	8.90

In conclusion, all formulations evaluated against naturally occurring populations of *An. albimanus* larvae in the Comayagua Valley of Honduras provided significant control ($P \leq 0.05$) within 24 hr and continued to provide such control through 240 hr posttreatment. Results from this study indicate that these formulations (Teknar, Arosurf MSF, Teknar + Arosurf, and Cuplex), costs notwithstanding, offer excellent potential as biorational alternative control strategies against this important malaria vector in Honduras.

REFERENCES

- Breeland, S.G. 1972. Studies on the ecology of *Anopheles albimanus*. *Am. J. Trop. Med. Hyg.* 21:751–754.
- Breeland, S.G. 1974. Population patterns of *Anopheles albimanus* and their significance to malaria abatement in El Salvador. *W.H.O. Symp. Malaria Res.* 73.23, 14 pp.
- Brown, A.W.A. 1986. Insecticide resistance in mosquitoes: A pragmatic review. *J. Am. Mosq. Control Assoc.* 2:123–140.
- Clyde, D.F. 1987. Recent trends in the epidemiology and control of malaria. *Epidemiol. Rev.* 9:219–243.
- Del Carmen, M., Sosa, E. and Bisset, J.A. 1984. Seasonal study of larval density of *Anopheles albimanus* (Wiedemann 1921) and some climatic and physio-chemical factors in an urban breeding place. *Rev. Cubana Med. Trop.* 36:288–296.
- Duncan, D.B. 1955. Multiple range test and multiple F tests. *Biometrics* 11:1–42.
- Elliott, R. 1969. Ecology and behavior of malaria vectors in the American region. *Cahiers ORSTOM Entomol. Med. Parasitol.* 7:29–33.
- Levy, R., Chlzzonlte, J.J., Garrett, W.D. and Miller, Jr., T.W. 1981. Ground and aerial application of a monomolecular organic surface film to control salt-marsh mosquitoes in natural habitats of southwestern Florida. *Mosq. News* 41:291–301.
- Levy, R., Powell, C.M., Hertlein, B.C. and Miller, Jr., T.W. 1984. Efficacy of Arosurf[®] MSF (monomolecular surface film) base formulations of *Bacillus thuringiensis* var. *israelensis* against mixed populations of mosquito larvae and pupae: Bioassay and preliminary field evaluations. *Mosq. News* 44:537–543.

- Margallt, J. and Bobroglio, H. 1984. The effect of organic materials and solids in water on the persistence of *Bacillus thuringiensis* var. *israelensis* serotype H-14. *Z. Angew. Entomol.* 97:516-520.
- Margallt, J. and Dean, D. 1985. The story of *Bacillus thuringiensis* var. *israelensis* (B.t.i.). *J. Am. Mosq. Control Assoc.* 1:1-7.
- Matsumura, F. 1975. *Toxicology of Insecticides*. Plenum Press, New York. 503 pp.
- Ohana, B., Margallt, J. and Barak, Z.E. 1987. Fate of *Bacillus thuringiensis* subsp. *israelensis* under simulated field conditions. *Appl. Environ. Microbiol.* 53:828-831.
- Pan American Health Organization. 1986. Malaria control in the Americas: A critical analysis. *Bull. PAHO* 20:17pp.
- Perlich, M.J., Rogers, J.T. and Boobar, L.R. 1987. Efficacy of Arosurf[®] MSF and formulations of *Bacillus thuringiensis* var. *israelensis* against *Anopheles albimanus*: Laboratory bioassay. *J. Am. Mosq. Control Assoc.* 3:485-488.
- Perlich, M.J., Rogers, J.T., Boobar, L.R. and Nelson, J.H. 1988. Laboratory evaluation of formulations of *Bacillus thuringiensis* var. *israelensis* combined with methoprene or a monomolecular surface film against *Anopheles albimanus* and *Anopheles stephensi*. *J. Am. Mosq. Control Assoc.* 4: (in press).
- Ramoska, W.A., Watts, S. and Rodriguez, R.E. 1982. Influence of suspended particulates on the activity of *Bacillus thuringiensis* serotype H-14 against mosquito larvae. *J. Econ. Entomol.* 75:1-4.
- Rozeboom, L.E. 1941. Distribution and ecology of the *Anopheles* mosquitoes of the Caribbean region. *Am. Assoc. Adv. Sci.* 15:98-107.
- SAS/STAT. 1985. *Guide for Personal Computers*. 6th ed. SAS Institute Inc., Cary, NC.
- Sloff, R. 1987. The control of malaria vectors in the context of The Health for All by the Year 2000 Global Strategy. *J. Am. Mosq. Control Assoc.* 3:551-555.
- van Essen, F. and Hembree, S. 1982. Simulated field studies with four formulations of *Bacillus thuringiensis* var. *israelensis* against mosquitoes: Residual activity and effect of soil constituents. *Mosq. News* 41:66-72.