

**THE EFFECT OF *PHENACOCCLUS MADEIRENSIS* GREEN (COCCOIDEA: PSEUDOCOCCIDAE) ON SOME BIOLOGICAL PARAMETERS OF FOUR SPECIES OF PHYTOSEIID MITES (PARASITIFORMES: PHYTOSEIIDAE)<sup>1</sup>**

S. RAGUSA DI CHIARA AND H. TSOLAKIS

*Istituto di Entomologia Agraria, Viale delle Scienze, 90128 Palermo, Italy*

**ABSTRACT**

The mealybug *Phenacoccus madeirensis* Green was supplied as prey to the following species of phytoseiid mites: *Euseius stipulatus* (Athias-Henriot), *Iphiseius degenerans* (Berlese), *Typhlodromus exhilaratus* Ragusa and *Typhlodromus cyptus* Athias-Henriot in order to assay its effect on their postembryonic development and oviposition rate. No young stages of the four species developed beyond the protonymph stage. Only very few eggs were laid by females of the four species. A list is provided with the results obtained while using other scale insect species, to show their importance for the survival of predaceous mites in the absence of optimal food.

**KEY WORDS:** Phytoseiidae, Pseudococcidae, postembryonic development, oviposition rate.

**INTRODUCTION**

Phytoseiid mites are important biological control agents of phytophagous mites. Many examples show the successes achieved using these predaceous mites in the field and greenhouses to control different phytophagous mites (McMurtry, 1982).

It is also known that most of them are facultative predators, which may develop and reproduce with success on various food substances (McMurtry and Scriven, 1966; Swirski et al., 1967a, b; Ragusa and Swirski, 1975; Pu et al., 1991), among which insects play also an important role. Of particular interest is the association of phytoseiid mites with scale insects.

During a survey on phytoseiid mites carried out in the surroundings of Palermo (Sicily) in May-July 1993, we found many specimens of *Iphiseius degenerans* (Berlese) and *Euseius stipulatus* (Athias-Henriot) associated with *Phenacoccus madeirensis* Green (Coccoidea: Pseudococcidae) living on *Eiythrina lantania* and *Acanthus* sp. Therefore it was decided to study the effect of the above-mentioned species on this mealybug. Moreover, as *Ph. madeirensis* also lives on citrus, we checked its influence on two other species of phytoseiid mites commonly found on citrus trees, namely *Typhlodromus exhilaratus* Ragusa and *T. cryptus* Athias-Henriot.

<sup>1</sup>Work supported with funds by Ministero dell'Universita e della Ricerca Scientifica e Tecnologica (40%).

### MATERIALS AND METHODS

The mites *I. degenerans*, *E. stipulatus*, *T. exhilaratus* and *T. cryptus* were collected in the field and reared in the laboratory. Cages used for mass breeding were described by Swirski et al. (1967a); a mixture of pollen of *Carpobrotus edulis*, *Malephora* sp. and *Oxalis cernua* was supplied as food. Five eggs per cage (5 replicates) were used to determine the influence on postembryonic development; oviposition was checked observing 3 young females and 1 male per cage (5 replicates) for a period of 10 days; if males died, they were immediately substituted with other young males in order to provide the full potential for copulation and oviposition. Eggs and crawlers of *Ph. madeirensis* were field collected and supplied as needed.

Trials were carried out in an air conditioned room at  $25 \pm 1^\circ\text{C}$  and  $75 \pm 5\%$  RH. Observations were conducted daily almost at the same hour.

### RESULTS AND DISCUSSION

The stages of *Ph. madeirensis* supplied had, in all cases, a negative influence on the postembryonic development (Table 1), as no immatures of the four mite species attained adulthood. Only larvae, which molted to protonymphs and died within approximately four days, were obtained. *E. stipulatus* had the highest larval mortality (88%) and *T. cryptus* the lowest (8%). Even if larvae and protonymphs were seen moving around the food and grasping it with their legs, we did not observe any predation. Larvae of *T. exhilaratus* do not feed (Ragusa, 1979), and probably the same happens with the other larvae, but further observations are required. A cannibalistic behavior was observed for *E. stipulatus*. All immatures were in poor conditions, thin, and mostly found on the surrounding cotton wool supplying water.

TABLE 1  
Effect of eggs and crawlers of *Phenacoccus madeirensis* Green on postembryonic development of *Typhlodromus exhilaratus* Ragusa, *Typhlodromus cryptus* Athias-Henriot, *Euseius stipulatus* (Athias-Henriot) and *Iphiseius degenerans* (Berlese)

Phytoseiids	Initial number of eggs	Mortality (%)				Attaining adulthood	Duration of protonymphal stage (days)
		Eggs	Larvae	Proto-nymphs	Deuto-nymphs		
<i>T. exhilaratus</i>	25	0	16	84	0	0	3.7
<i>T. cryptus</i>	25	0	8	92	0	0	4.2
<i>E. stipulatus</i>	25	0	88	12	0	0	3.8
<i>I. degenerans</i>	25	0	43	57	0	0	3.6

The value of the food supplied for oviposition proved to be very low (Table 2). The highest oviposition was obtained with *T. cryptus* (0.38 eggs/female/day), but it was almost nil for the other species. Eggs were laid between the second and the fourth day.

No adults of *E. stipulatus* and *I. degenerans* survived at the end of the experiment; mortality was low for *T. exhilaratus* (20%) and higher for *T. cryptus* (46.6%).

TABLE 2

Effect of eggs and crawlers of *Phenacoccus madeirensis* Green on the oviposition rate and survival of *Typhlodromus exhilaratus* Ragusa, *Typhlodromus cryptus* Athias-Henriot *Euseius stipulatus* (Athias-Henriot) and *Iphiseius degenerans* (Berlese)

Phytoseiids	Number of adults		Number of adults surviving at the end of the test		Mortality		Eggs/female/day
	Females	Males	Females	Males	Females	Males	
<i>T. exhilaratus</i>	15	5	12	1	20	80	0.11
<i>T. cryptus</i>	15	5	8	1	46.6	80	0.38
<i>E. stipulatus</i>	15	5	0	0	100	100	0.10
<i>I. degenerans</i>	15	5	0	0	100	100	0.08

It is known that some mites feed upon scale insects: species belonging to the genus *Hemisarcoptes* have been used in the biological control of armored scale insects (Gerson et al., 1990). Studies were conducted to evaluate the impact of *Hemisarcoptes* on host scales (Gerson and Schneider, 1981). More recently the potential of *Hemisarcoptes coccophagus* Meyer in controlling the kiwi pest *Hemiberlesia lataniae* (Signoret) was shown after its introduction into New Zealand (Hill et al., 1993).

As far as phytoseiid mites are concerned, many studies were carried out to assay the influence of various stages of different scale insects on their postembryonic development and oviposition rate.

In Table 3 we summarized results obtained by various authors using different species of phytoseiid mites. The results are controversial as the influence was sometimes positive and in other cases negative; when California red scale *Aonidiella aurantii* (Maskell) was used, the influence on postembryonic development was positive and the percentages of adults obtained ranged from 38 to 100%. Immatures of *Amblyseius limonicus* (Garman and McGregor) feeding on crawlers and honeydew of *Pseudococcus* aff. *citriculus* Green reached adulthood in 26.3% cases. Only 16% immatures of *Amblyseius andersoni* (Chant) fed crawlers of *Saissetia oleae* (Olivier) reached adulthood. No other scale insects had a positive influence on the postembryonic development: usually it was nil or very poor (0–4.8%).

The influence of scale insects on oviposition was similar to that on postembryonic development; this was the case with *Aonidiella aurantii*. Sometimes the effect was more positive on oviposition rate than on postembryonic development. This was the case of *Amblyseius swirskii* (Athias-Henriot) which laid 1.11 eggs/female/day, while 38.2% of its immatures reached adulthood. Generally speaking, other scales such as *Ceroplastes floridensis* Comstock, *Pseudococcus* aff. *citriculus*, *Coccus hesperidum* Linnaeus, *Saissetia oleae* (Olivier), *S. coffeae* (Walker), *Pseudococcus longispinus* (Targioni Tozzetti), *Icerya purchasi* Maskell and *Phenacoccus madeirensis* did not influence oviposition positively, as the number of eggs was almost nil.

Muma (1971) reviewed the food habits of phytoseiid mites, classifying foods in four categories in relation to the influence they had on postembryonic development and oviposition rate, as follows: (1) optimal, (2) adequate, (3) survival, (4) inadequate. Using this classification,

TABLE 3  
Some reported data on the postembryonic development and oviposition rate of various species of phytoseiid mites feeding on different scale insects under laboratory conditions

Phytoseiids	Hosts	Attaining adulthood (%)	Eggs/female/day	References
<i>T. rickeri</i>	<i>Hemiberlesia lataniae</i>	—	low	McMurtry, 1963
<i>A. limonicus</i>	<i>Hemiberlesia lataniae</i>	—	0.6	McMurtry, 1963
	<i>Aonidiella aurantii</i>	85.7	—	Swirski and Dorzia, 1968
<i>E. hibisci</i>	<i>Pseudococcus</i> aff. <i>citriculus</i>	26.3	0.23	Swirski and Dorzia, 1968
	<i>Hemiberlesia lataniae</i>	—	0.4	McMurtry, 1963
<i>A. chilensis</i>	<i>Aonidiella aurantii</i>	78.2	0.7	Swirski et al., 1970
	<i>Ceroplastes floridensis</i>	0	—	Swirski et al., 1970
<i>Metaseiulus occidentalis</i>	<i>Aonidiella aurantii</i>	88.9	0.53	Swirski et al., 1970
	<i>Ceroplastes floridensis</i>	0	0	Swirski and Dorzia, 1969
<i>T. athiasae</i>	<i>Aonidiella aurantii</i>	79.2	0.57	Swirski and Dorzia, 1969
	<i>Hemiberlesia lataniae</i>	—	0.25	McMurtry, 1963
<i>T. athiasae</i>	<i>Aonidiella aurantii</i>	100	0.86	Swirski et al., 1967b
	<i>Pseudococcus</i> aff. <i>citriculus</i>	0	—	Swirski et al., 1967b

<i>A. swirskii</i>	<i>Aonidiella aurantii</i>	crawlers	38.2	1.11	Swirski et al., 1967a	
	<i>Pseudococcus aff. citriculus</i>	crawlers and honeydew	0	0.01	Swirski et al., 1967a	
	<i>Coccus hesperidum</i>	crawlers	0	0	Ragusa and Swirski, 1977	
	<i>Coccus hesperidum</i>	crawlers and honeydew	0	0.02	Ragusa and Swirski, 1977	
	<i>Saissetia oleae</i>	crawlers	0	0	Ragusa and Swirski, 1977	
	<i>Saissetia oleae</i>	eggs and crawlers	4.8	0	Ragusa and Swirski, 1977	
	<i>Saissetia oleae</i>	crawlers and honeydew	4.2	0	Ragusa and Swirski, 1977	
	<i>Saissetia oleae</i>	eggs, crawlers and honeydew	0	0	Ragusa and Swirski, 1977	
	<i>Saissetia coffeae</i>	eggs	0	0	Ragusa and Swirski, 1977	
	<i>Saissetia coffeae</i>	eggs, crawlers and honeydew	0	0	Ragusa and Swirski, 1977	
	<i>Pseudococcus aff. citriculus</i>	crawlers and honeydew	0	0	Ragusa and Swirski, 1977	
	<i>Pseudococcus longispinus</i>	crawlers and honeydew	4.2	0	Ragusa and Swirski, 1977	
	<i>E. rubini</i>	<i>Aonidiella aurantii</i>	crawlers	71.4	0.48	Swirski et al., 1967a
		<i>Pseudococcus aff. citriculus</i>	crawlers and honeydew	0	0.13	Swirski et al., 1967a
<i>A. andersoni</i>	<i>Saissetia oleae</i>	eggs and crawlers	16	0.01	Ragusa Di Chiara and Tsolakis, 1994	
	<i>Icerya purchasi</i>	eggs and crawlers	0	0	Ragusa Di Chiara and Tsolakis, 1994	
<i>T. exthilaratus</i>	<i>Phenacoccus madeirensis</i>	eggs and crawlers	0	0.11	Present paper	
	<i>Phenacoccus madeirensis</i>	eggs and crawlers	0	0.38	Present paper	
<i>E. stipulatus</i>	<i>Phenacoccus madeirensis</i>	eggs and crawlers	0	0.1	Present paper	
<i>I. degenerans</i>	<i>Phenacoccus madeirensis</i>	eggs and crawlers	0	0.08	Present paper	

we may consider most scales as either adequate or survival foods, and few of them as inadequate. But even if the nutritional value of scale insects is inconclusive, they can still be used by phytoseiid mites during periods when the optimal food is absent to maintain a population which will increase as soon as better nutrients appear again. Moreover, phytoseiid mites can also find overwintering shelter under scale covers. Such behavior explains why many mites are often found under scale covers even if the species has no positive influence on the predaceous mites. These could account for the cases reported by Putman (1962) and by Lal and Naji (1980); the latter found 10–15% of scales with phytoseiid mites during winter.

In conclusion, the reason why we found many specimens of *E. stipulatus* and *I. degenerans* associated with *Ph. madeirensis* during May–July must still be elucidated.

#### REFERENCES

- Gerson, U. and Schneider, R.** 1981. Laboratory and field studies on the mite *Hemisarcoptes coccophagus* Meyer (Astigmata: Hemisarcoptidae), a natural enemy of armored scale insects. *Acarologia* 22:199–208.
- Gerson, U., O'Connor, B.A. and Houck, M.A.** 1990. Acari. In: Armored Scale Insects: Their Biology, Natural Enemies and Control. Vol. 4B. Edit. D. Rosen. Elsevier, Amsterdam. pp. 77–97.
- Hill, M.G., Allan, D.J., Henderson, R.C. and Charles, J.C.** 1993. Introduction of armored scale predators and establishment of the predatory mite *Hemisarcoptes coccophagus* (Acari: Hemisarcoptidae) on latania scale, *Hemiberlesia latania* (Homoptera: Diaspididae) in kiwifruit shelter trees in New Zealand. *Bulletin of Entomological Research* 83:369–376.
- Lal, O.P. and Naji, A.H.** 1980. Observations on the predators of the black olive scale, *Saissetia oleae* Bern. (Homoptera: Coccidae) in the Socialist Peoples Libyan Arab Jamahiriya. *Journal of Plant Diseases and Protection* 81:27–31.
- McMurtry, J.A.** 1963. Diaspine scale insects as prey for certain phytoseiid mites. In: Advances in Acarology. Vol. 1. Edit. J.A. Naegele. Cornell University Press, Ithaca, N.Y. pp. 151–154.
- McMurtry, J.A.** 1982. The use of phytoseiids for biological control: Progress and future prospects. In: Recent Advances in Knowledge of the Phytoseiidae. Edit. M.A. Hoy. University of California, Publication No. 3284. pp. 23–48.
- McMurtry, J.A. and Scriven, G.T.** 1966. The influence of pollen and prey density on the number of prey consumed by *Amblyseius hibisci* (Acarina: Phytoseiidae). *Annals of the Entomological Society of America* 59:147–149.
- Muma, H.M.** 1971. Food habits of Phytoseiidae (Acarina: Mesostigmata) including common species on Florida citrus. *The Florida Entomologist* 54:21–34.
- Pu, T., Zeng, T. and Wei, D.** 1991. An integrated evaluation on 20 plant pollen as diet for mass rearing *Amblyseius makuwa* (Acari: Phytoseiidae). *Chinese Journal of Biological Control* 7: 111–114.
- Putman, W.L.** 1962. Life-history and behaviour of the predaceous mite *Typhlodromus (T.) caudiglans* Schuster (Acarina: Phytoseiidae) in Ontario, with notes on the prey of related species. *Canadian Entomologist* 94:163–177.
- Ragusa, S.** 1979. Laboratory studies on the food habits of the predaceous mite *Typhlodromus exhilaratus*. *Recent Advances in Acarology* 1:485–490.
- Ragusa, S. and Swirski, E.** 1975. Feeding habits, development and oviposition of the predaceous mite *Amblyseius swirskii* Athias-Henriot (Acarina: Phytoseiidae) on pollen of various weeds. *Israel Journal of Entomology* 10:93–103.
- Ragusa, S. and Swirski, E.** 1977. Feeding habits, post-embryonic and adult survival, mating, virility and fecundity of the predaceous mite *Amblyseius swirskii* (Acarina: Phytoseiidae) on some coccids and mealybugs. *Entomophaga* 22:383–392.

- Ragusa Di Chiara, S. and Tsolakis, H.** 1994. Influence of different kinds of food substances on the postembryonic development and oviposition rate of *Amblyseius andersoni* (Chant) (Parasitiformes, Phytoseiidae). II<sup>nd</sup> Symposium EURAAC, Krynica 1992, Poland (in press).
- Swirski, E., Amitai, S. and Dorzia, N.** 1967a. Laboratory studies on the feeding, development and reproduction of the predaceous mites *Amblyseius rubini* Swirski and Amitai and *Amblyseius swirskii* Athias (Acarina: Phytoseiidae) on various kinds of food substances. *Israel Journal of Agricultural Research* 17:101–119.
- Swirski, E., Amitai, S. and Dorzia, N.** 1967b. Laboratory studies on the feeding, development and oviposition of the predaceous mite *Typhlodromus athiasae* P. and S. (Acarina: Phytoseiidae) on various kinds of food substances. *Israel Journal of Agricultural Research* 17:213–218.
- Swirski, E., Amitai, S. and Dorzia, N.** 1970. Laboratory studies on the feeding habits, post-embryonic survival and oviposition of the predaceous mites *Amblyseius chilensis* Dosse and *Amblyseius hibisci* Chant (Acarina: Phytoseiidae) on various kinds of food substances. *Entomophaga* 15:93–106.
- Swirski, E. and Dorzia, N.** 1968. Studies on the feeding, development and oviposition of the predaceous mite *Amblyseius limonicus* Garman and McGregor (Acarina: Phytoseiidae) on various kinds of food substances. *Israel Journal of Agricultural Research* 18:71–75.
- Swirski, E. and Dorzia, N.** 1969. Laboratory studies on the feeding, development and fecundity of the predaceous mite *Typhlodromus occidentalis* Nesbitt (Acarina: Phytoseiidae) on various kinds of food substances. *Israel Journal of Agricultural Research* 19:143–145.